

References

- Alkaade, S., & Vareedayah, A. A. (2014). Managing high-output stomas. *Gastroenterology & Hepatology*, 10(11), pp. 756–763.
- Baker, M. L., Williams, R. N., & Nightingale, J. M. (2011). Causes and management of a high-output stoma. *Colorectal Disease*, 13(2), pp. 191–197.
- Black, P., & Cottrell, S. (2019). Stoma care: Multidisciplinary approaches to improving patient outcomes. *British Journal of Nursing*, 28(Sup14), pp. S4–S12.
- BMJ (2020). Management of dehydration in adults. *BMJ*, 370, m2780. [online] Available at: <https://www.bmj.com/content/370/bmj.m2780> [Accessed 10 Jan. 2025].
- British Dietetic Association (BDA) (2022). Nutritional management of high-output stomas: Best practice guidelines.
- Brown, H., & Randle, J. (2005). Living with a stoma: A review of the literature. *Journal of Clinical Nursing*, 14(1), pp. 74–81.
- Cameron, I., et al. (2010). High-output stomas: Guidelines for management. *British Journal of Nursing*, 19(17), pp. 1062–1069.
- Davenport, R., Mackenzie, K., & Stewart, J. (2013). Management of peristomal skin complications in high-output stoma patients. *Journal of Wound Care*, 22(12), pp. 692–700.
- ESPEN Guidelines (2018). Clinical nutrition and enteral feeding best practices.
- Gabe, S. M., et al. (2005). Managing high-output stomas and enterocutaneous fistulas. *Frontline Gastroenterology*, 6(3), pp. 118–124.
- Holman, N., & Flook, R. (2020). Guidelines for the management of high-output stomas. *British Journal of Nursing*, 29(20), pp. S14–S20.
- Johnson, P., & Hall, S. (2018). Personalized care planning in chronic illness. *Nursing Times*, 114(3), pp. 26–29.
- McEvoy, P., & Duffy, A. (2008). Holistic practice – A concept analysis. *Nurse Education in Practice*, 8(6), pp. 412–419.
- Mount, D. (2015). Hyponatremia: Evaluation and management. *The New England Journal of Medicine*, 372(1), pp. 55–65.
- NICE Guidelines (2021). Enteral and parenteral nutrition in clinical practice.
- NICE Guidelines (2021). Stoma care: Management and dietetic considerations.
- Nightingale, J. M., & Lennard-Jones, J. E. (1999). Nutritional assessment and intestinal failure. *Nutrition*, 15(4), pp. 302–307.
- Nightingale, J. M., & Woodward, J. M. (2006). Guidelines for management of patients with a high-output stoma. *Gut*, 55(Suppl 4), pp. iv1–iv12.
- Nightingale, J. M., & Woodward, J. M. (2006). Management of patients requiring parenteral nutrition. *Gut*, 55(Suppl 4), pp. iv1–iv12.
- Nightingale, J. M. (2003). Management of patients with a short bowel. *Postgraduate Medical Journal*, 79(937), pp. 178–183.
- Persson, E., Hellström, A. L., & Hallén, A. M. (2010). Experiences of living with a permanent stoma. *Ostomy Wound Management*, 56(6), pp. 20–27.
- Pironi, L., et al. (2016). ESPEN guidelines on chronic intestinal failure. *Clinical Nutrition*, 35(2), pp. 247–307.
- Pironi, L., Arends, J., Baxter, J., et al. (2020). Management of high-output enterostomies. *Clinical Nutrition*, 39(4), pp. 1181–1193.
- Powell-Tuck, J., et al. (2008). Fluid and electrolyte balance in clinical practice. Hodder Arnold.
- Smith, G., Finch, T., & Shannon, R. (2017). Empowering patients in stoma care. *British Journal of Nursing*, 26(22), pp. S20–S25.
- Wade, B. E. (2016). Psychological impact of stoma surgery. *Gastrointestinal Nursing*, 14(6), pp. 21–27.
- World Health Organization (2017). Oral rehydration salts: Production of the new ORS. Geneva: WHO. Available at: <https://www.who.int/publications/i/item/WHO-FCH-CAH-06.1> [Accessed 10 Jan. 2025].